


FUNCTIONAL NASAL SURGERY POST OPERATIVE INSTRUCTIONS



Norwest ENT
Group

Leaving the hospital

1. If you have not already been given a postoperative appointment, please ring and arrange a visit for approximately 4 weeks after surgery.
2. Any medication you are prescribed, should be taken as directed on the bottle.
3. There are no dietary restrictions, although after a general anaesthetic it may be best to start with clear liquids.

Nasal Packing

Most nose operations do not require nasal packing that needs to be removed after surgery. Occasionally if packs are placed in the nose after surgery, then strings will be tied in front of the nose. The packs will then be removed before you go home.

Nasal Washing

Nasal douching (washing) is recommended after all nasal surgery for 3 months. Commence douching 2 days after the operation. Use the pump sprays or nasal douches (FLO, Sinus Rinse) 4 to 6 times per day. Vaseline around the nostril following nasal douching will reduce discomfort and skin irritation.

Precautions

1. GENTLE nose blowing is OK 48 hours after surgery. Any accumulated secretions in the back of the nose may be drawn back into the throat and expectorated if desired. This is particularly important if you develop a cold.
2. If it is necessary to sneeze, do so with your mouth open.
3. DO NOT take any unnecessary chance of catching a cold. Avoid undue exposure or fatigue. Should you catch a cold, treat it in your usual way.
4. You may anticipate a certain amount of discomfort after the surgery, but this is typically well controlled by pain tablets.
5. DO NOT plan to drive a car home from the hospital. Someone should drive you home.
6. DO NOT perform any heavy lifting (more than 15 kilograms) or vigorous physical activity for three weeks after surgery.
7. Some nose bleeding is expected after the surgery. To control this bleeding, sit up, bend the head forward and pinch the nose closed with thumb and forefinger. Holding this position for 10 minutes by the clock will settle most bleeding down. Please contact your doctor if the bleeding does not stop.

Pain Management

- Paracetamol (panadol) 500mg tablets 2 tablets four times a day.
- Oxycodone tablets when required and as prescribed.
- If these measures are not controlling your pain then contact your anaesthetist or us at the office.

Nausea and Vomiting

While it is not uncommon to get some nausea and or vomiting after a nasal surgery it is not usually severe or long lasting. If you find you have persistent nausea and vomiting then have the ondansetron prescription filled. These are small wafers that you dissolve under your tongue. If after 24 hours of taking the ondansetron you are unable to keep fluids down you should contact your anaesthetist or Dr Patel as you may then require intravenous fluids.

Contact Numbers:

Dr Nirmal Patel
M 0431 444 636

Dr Tobias Pincock
M 0467 300 050

Dr Leo Pang
M 0414 878 398